

## *Salmonella* Surveillance Among Great-Tailed Grackles (*Quiscalus mexicanus*) and Other Urban Bird Species in Eastern Texas

Mary K. Grigar,<sup>1</sup> Kevin J. Cummings,<sup>1</sup> Lorraine D. Rodriguez-Rivera,<sup>1</sup> Shelley C. Rankin,<sup>2</sup> Krista Johns,<sup>1</sup> Gabriel L. Hamer,<sup>3</sup> and Sarah A. Hamer<sup>1</sup>

### Abstract

Wild birds may play an important role in maintaining and transmitting *Salmonella*. Their ability to travel large distances and their proximity to human habitations could make them a vehicle for bridging *Salmonella* from wild and domestic animals to humans. To determine the potential public health risk presented by urban birds, we investigated the prevalence of *Salmonella* among great-tailed grackles (*Quiscalus mexicanus*) and other cohabiting urban bird species. Fecal samples were collected from 114 birds communally roosting in parking lots of retail locations in Brazos County, Texas, from February through July of 2015. Great-tailed grackles and European starlings (*Sturnus vulgaris*) were the predominant species sampled. Standard bacteriologic culture methods were used to isolate *Salmonella* from samples, and isolates were characterized by serotyping and antimicrobial susceptibility testing. Overall, 1.8% (2/114) of samples were confirmed positive for *Salmonella*. Both positive birds were great-tailed grackles sampled in June, yielding a 2.6% (2/76) apparent prevalence among this species. Isolates were serotyped as *Salmonella* Typhimurium and found to be pan-susceptible based on the National Antimicrobial Resistance Monitoring System (NARMS) panel of antimicrobial agents. The occurrence of *Salmonella* in great-tailed grackles represents a potential threat to public health, particularly considering their population size and tendency to congregate near human establishments such as grocery stores.

**Keywords:** birds, epidemiology, *Salmonella*, surveillance, zoonotic

### Introduction

**S**ALMONELLA ENTERICA IS a zoonotic pathogen that poses a considerable threat to public health. Transmission occurs through foodborne exposure as well as direct contact with the feces of infected animals. *Salmonella* is estimated to cause ~1.2 million illnesses, 20,000 hospitalizations, and 400 deaths annually in the United States (Scallan et al. 2011), at an economic cost estimated to be \$4.4 billion per year (Scharff 2012). Typical disease manifestations in humans include diarrhea, fever, anorexia, abdominal pain, vomiting, and malaise (Heymann 2015). Livestock and poultry species serve as key reservoirs for *Salmonella*, although the organism has been isolated from the gastrointestinal tract of a wide range of wild and domestic host species (Skov et al. 2008). However, the epidemiology of *Salmonella* among wildlife has been poorly defined, relative to domestic species.

Wild birds may play a critical role in disseminating *Salmonella* and other enteric pathogens across the landscape, given their ability to travel great distances during migrations, dispersals, or between feeding and roosting activities. Thus, wild birds have the potential to pose a substantial risk to public health (Callaway et al. 2014). To this end, there have been several studies involving birds near agricultural operations such as dairies and feedlots. For example, Callaway et al. (2014) conducted a survey of birds often associated with cattle in Brazos County, Texas, namely brown-headed cowbirds (*Molothrus ater*), common grackles (*Quiscalus quiscula*), and cattle egrets (*Bubulcus ibis*), and found 14.9% of these birds to be shedding *Salmonella*. Phalen et al. (2010) estimated a *Salmonella* prevalence ranging from 29% to 91% among cattle egret nestlings from various colonies in central Texas. The serotypes identified in these birds were also isolated from horses in the 2 years following their study,

<sup>1</sup>Department of Veterinary Integrative Biosciences, College of Veterinary Medicine & Biomedical Sciences, Texas A&M University, College Station, Texas.

<sup>2</sup>School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania.

<sup>3</sup>Department of Entomology, College of Agriculture and Life Sciences, Texas A&M University, College Station, Texas.

indicating that egrets and horses may be exposed to similar sources of *Salmonella* or that transmission may be occurring between them. In addition, Kirk et al. (2002) captured 892 birds at nine dairies in California and isolated *Salmonella* spp. from 1.2% to 3.2% of each avian species, with brown-headed cowbirds and house sparrows (*Passer domesticus*) having the highest prevalence. Fewer studies have been conducted in nonagricultural settings. In particular, there is a dearth of research on *Salmonella* shedding among bird species that congregate in urban locations with high human activity, such as great-tailed grackles (*Quiscalus mexicanus*). Hamer et al. (2012) collected fecal samples from 180 birds in the greater Chicago, IL area and found only one individual (0.6%) to be infected with *Salmonella*. Janecko et al. (2015) identified *Salmonella* in 1.7% (10/590) of samples from American crows collected in four states.

Great-tailed grackles spread from Mexico to reach southern Texas by the 1880s, and their range now extends as far north and west as South Dakota, Idaho, and California (Wehtje 2003). In the United States, the species is highly adapted to the urban environment and forms large aggregations of communally roosting individuals at night. The locations of these nocturnal roosts are frequently the trees of large, well-illuminated parking lots (Hall and Harvey 2007). The presence of heavy loads of avian feces beneath roost trees, coupled with raucous vocalizations, result in these birds and other urban bird species gaining a reputation as pests, with many establishments attempting to haze these birds away (Fitzwater et al. 1988). The objectives of this study were to estimate the prevalence of *Salmonella* shedding among roosting great-tailed grackles and other urban birds in the parking lots of retail locations in Brazos County, Texas, and characterize the isolates through serotyping and antimicrobial susceptibility testing.

## Materials and Methods


### Bird collection and processing

On nine evenings from February through July 2015, samples were collected from roosting birds in urban parking lots of retail buildings in College Station, Brazos County, located in eastern Texas. Great-tailed grackles have been established since the 1940s in many metropolitan areas of Texas, including

College Station (Wehtje 2003). Sites were selected based on prior observations of Great-tailed Grackle and other urban bird aggregations for staging and roosting. The study area was ~9 hectares in size and encompassed two gas stations, a bank, a grocery store, and several restaurants and other businesses. Nylon mist nets (12-meter length; Association of Field Ornithologists, Portland, ME) were erected next to communal roost trees as previously described (Hamer et al. 2012). As roost tree canopies were above the standard height of a mist net, we adapted this protocol by connecting two rigid conduit pipes with a threaded couplet to create 6.1-meter poles that supported the nets (Fig. 1A). Mist nets with 30 mm mesh size were used initially, but larger birds (e.g., male great-tailed grackles) were observed escaping from these nets; thus, we switched to 61 mm mesh size nets to improve capture success. The nets were erected 1–4 h before sunset as the communal birds were staging. Birds were captured in nets as they arrived at the roost trees and when 3-meter conduit poles were used to flush roosting birds into the nets. Birds were extracted immediately after being caught in many cases, and nets were monitored for additional captured birds every 15–20 min. A double layer of new paper bags was used to hold birds individually until they were ready to be processed.

For each captured bird, the species, weight, wing chord, tail length, sex, and age class (hatch year or after hatch year) were recorded. Sterile cotton-tipped swabs were used to collect fecal samples from the inner paper bag holding the bird, directly from the cloaca, or in a few cases from processing tables or technicians; samples were then deposited into Amies media (Becton-Dickinson, Sparks, MD). When voided fecal samples were available, we attempted to collect the entire amount. Standard procedures were utilized to prevent cross-contamination of samples, such as changing gloves between handling individual birds. Up to two samples were collected per bird and kept in a cooler at ~4°C. Following sample collection, birds were released at the site of capture after attaching a United States Fish and Wildlife Service (USFWS) metal leg band with a unique number, or in some cases, they were euthanized for unrelated research efforts. Fieldwork was conducted with approvals from the Institutional Animal Care and Use Committee at Texas A&M University, U.S. Fish and Wildlife Service, U.S. Geological Survey, Texas Parks and Wildlife Department, and private business owners.

**FIG. 1.** Nylon mist nets on 6.1-meter poles next to a roost tree in the foreground with staging great-tailed grackles (*Quiscalus mexicanus*) on power lines in the background (A) and white feces from great-tailed grackles under two roost trees in a parking lot in College Station, Texas (B). A female great-tailed grackle on grocery store shopping carts (C) and avian feces on the handle of a shopping cart (D). Color images available online at [www.liebertpub.com/vbz](http://www.liebertpub.com/vbz)



### Estimate of study population

Communal Great-tailed Grackle roosts are dynamic in the College Station, Texas, environment. The size and location of these roosts vary throughout the year, likely due to breeding activities and hazing practices to disrupt the large aggregations of birds. One technique to census communal roosting birds is to station multiple observers around the roost to count birds as they arrive (Rumbold et al. 2009). As great-tailed grackles arrive at their communal roost, however, they stage in large numbers and are prone to flight when disturbed. Mass departure from staging locations or roost trees occurs throughout dusk and into the evening, triggered by the many forms of disturbance that are common in parking lots of retail buildings in urban areas. To avoid duplicate counting of birds, we pursued an alternative method to estimate local population size. During the nine evenings of sampling (February through July 2015) and on frequent scouting visits before and during the sampling period, we became familiar with the trees utilized to roost by great-tailed grackles and the boundaries of the roost where trees were not used. Roost trees were primarily Southern live oaks (*Quercus virginiana*), with a typical height of ~8 meters, located in parking lots near artificial lighting. We estimated that ~55 roost trees were being utilized by great-tailed grackles in the study area on any given night, based on our experience in the field combined with satellite imagery. Dispersal of grackles from roost trees, in response to our sampling efforts or other human activity, allowed a conservative estimation of ~40 birds per tree. Thus, we estimated a total of 2,200 great-tailed grackles within the study area, utilizing the communal roosts (Fig. 1A–D).

### Microbiologic procedure for *Salmonella* detection

Within 1–5 h of collection, samples were brought to the research laboratory and enriched in 5 mL of tetrathionate (TT) broth (Becton, Dickinson and Company, Franklin Lakes, NJ) containing 0.1 mL of iodine solution. Enrichments were incubated at 42°C for 24 h and then plated to xylose lysine tergitol-4 (XLT-4) agar (Northeast Laboratories, Waterville, ME). Plates were incubated for 48 h at 37°C; these were examined for suspected colonies at 24 and 48 h. Following incubation, a single suspected colony was transferred to Kligler iron agar (KIA) slants (Becton, Dickinson and Company) and incubated for 18–24 h at 37°C. Colonies from positive slants were streaked onto trypticase soy agar (TSA) 5% sheep blood (Becton, Dickinson and Company) and incubated at 37°C for 18–24 h. After incubation, one positive colony was transferred to brain–heart infusion (BHI) broth (Becton, Dickinson and Company), incubated at 37°C for 16–24 h, and then frozen in a 15% glycerol solution at –80°C for further characterization.

Following the first month of sampling, samples were processed using the preceding method in parallel with an additional method based on Andrews and Hammack (2007). This second method was incorporated to allow future studies to characterize non-*Salmonella* bacteria that may be present in the samples and further improve *Salmonella* recovery. When two samples per bird were available, they were randomly allocated to each method. If only one sample was available, it was processed using the modified Andrews and Hammack (2007) method. This modified method consisted of

the same steps as previously described, but with the addition of a nonspecific enrichment step in tryptic soy broth (TSB; Becton, Dickinson and Company) before the TT enrichment and an enrichment in Rappaport-Vassiliadis (RV) broth (Becton, Dickinson and Company) following TT enrichment. TSB enrichment consisted of incubating the swabs for 2 h at room temperature followed by at least 10-h incubation at 37°C. One milliliter of TSB was transferred to 9 mL of TT broth and incubated for 24 h at 37°C. The remaining TSB was incubated for an additional 12 h and then frozen in the same manner as BHI in the previously described method. Then, 200 µL of the TT broth was transferred to 5 mL of RV broth and incubated for 24 h at 42°C. After incubation, RV was plated to XLT-4. Steps following the 48-h incubation are the same as in the previously described method. In addition, presumptive *Salmonella* isolates were confirmed by amplification and detection of the *invA* gene using PCR as previously described (Kim et al. 2007).

### Serotyping

Confirmed *Salmonella* isolates were sent to the Clinical Microbiology Laboratories at the University of Pennsylvania Veterinary Hospital for serotyping. The xMAP<sup>®</sup> *Salmonella* Serotyping Assay (Luminex, Austin, TX) was used to identify *Salmonella* isolates. The assay simultaneously determines O and H antigen genes and also identifies serotype-specific markers in the AT (Additional Targets) test. *Salmonella* template DNA was extracted using the InstaGene Matrix as described by the manufacturer (Bio-Rad, Hercules, CA). Multiplex PCR was performed, and the amplicons were hybridized with the oligonucleotide probe-coupled bead mixture and then labeled with streptavidin-R-phycoerythrin (SAPE)<sup>®</sup> reporter. The assay plate was analyzed on a Luminex<sup>®</sup> LX200<sup>™</sup> platform, and the data were exported to Excel (Microsoft, Redmond, WA) for analysis.

### Antimicrobial susceptibility testing

Antimicrobial susceptibility of confirmed *Salmonella* isolates was determined by the broth microdilution method. Minimal inhibitory concentrations (MIC) were established for each isolate against the National Antimicrobial Resistance Monitoring System (NARMS) Gram-negative panel of 14 antimicrobial agents (plate code CMV3AGNF, Sensititre; TREK Diagnostic Systems, Cleveland, OH): amoxicillin/clavulanic acid, ampicillin, azithromycin, cefoxitin, ceftiofur, ceftriaxone, chloramphenicol, ciprofloxacin, gentamicin, nalidixic acid, streptomycin, sulfisoxazole, tetracycline, and trimethoprim/sulfamethoxazole. Clinical and Laboratory Standards Institute (CLSI) guidelines were used to interpret MIC values when available (CLSI 2008, 2012). Otherwise, MIC values were interpreted using NARMS breakpoints (FDA 2013). Quality control was performed weekly using *Escherichia coli* ATCC 25922.

### Results

In total, 117 birds were captured, including 76 great-tailed grackles, 34 European starlings (*Sturnus vulgaris*), 4 house sparrows, 1 brown-headed cowbird, 1 cedar waxwing (*Bombycilla cedrorum*), and 1 Eurasian collared-dove (*Streptopelia decaocto*). The number of birds caught per

TABLE 1. DISTRIBUTION OF SEX, AGE, SEASON, AND STATUS OF *SALMONELLA* SHEDDING AMONG 117 URBAN BIRDS CAPTURED IN COLLEGE STATION, TEXAS, 2015

	<i>Great-tailed grackle</i>	<i>European starling</i>	<i>House sparrow</i>	<i>Brown-headed cowbird</i>	<i>Cedar waxwing</i>	<i>Eurasian collared-dove</i>	<i>Salmonella culture positive</i>
Sex							
Male	11	7	3	0	0	0	0
Female	59	3	0	1	0	0	2
Unknown	6	24	1	0	1	1	0
Age							
Hatch year	5	0	1	0	0	0	0
After hatch year	69	34	3	1	1	1	2
Unknown	2	0	0	0	0	0	0
Season <sup>a</sup>							
Spring	49	29	2	1	1	1	0
Summer	27	5	2	0	0	0	2
Total	76	34	4	1	1	1	2

<sup>a</sup>Spring includes February through April; summer includes May through July.

night ranged from 1 to 22, with a median of 16 birds. Of the birds captured, 63 were female, 21 were male, and 33 were of unknown sex. The majority of birds captured ( $n = 109$ ) were determined to have hatched before the calendar year of capture (Table 1). All birds were judged to be in adequate health condition based on gross appearance, with no overt signs of disease.

Fecal samples were collected from 114 birds, with 34 sampled solely from collection bags, 34 solely from cloacal swabs, 37 using both methods, 6 from other surfaces, and 3 with no record of collection method. Overall, 1.8% (2/114; 95% CI 0.2%–6.2%) of samples were confirmed positive for *Salmonella*. Both positive birds were great-tailed grackles sampled in June, yielding a 2.6% (2/76; 95% CI 0.3%–9.2%) apparent prevalence among this species. The great-tailed grackles were sampled at different roosting sites and by different methods (collection bag and cloacal swab). Both isolates were serotyped as *Salmonella* Typhimurium, and both were pan-susceptible to the antimicrobial agents included in the NARMS panel.

## Discussion

The occurrence of *Salmonella* in great-tailed grackles represents a potential threat to public health, particularly in view of their population size and preferred habitat. Great-tailed grackles tend to congregate near human establishments, including grocery stores, restaurants, and gas stations, creating opportunities for zoonotic transmission through direct contact or foodborne exposure. Assuming that our sample is representative of the population of great-tailed grackles in the study area, we estimate that over 50 *Salmonella*-positive birds were present in this area on a given night during the study period, based on the apparent prevalence among this species (2.6%) and our conservative estimate of local population size (2,200). This is likely to be an underestimate, as fecal culture does not have perfect sensitivity for detecting the presence of *Salmonella*.

In addition to detection methods, sampling location and time of year influenced our results. Studies on *Salmonella* shedding among wild birds in agricultural settings have

generated higher prevalence estimates (Callaway et al. 2014), suggesting that roosting near dairies and feedlots may be a risk factor for positive *Salmonella* status among birds. Birds in urban locations may have a lower risk of *Salmonella* exposure, relative to birds that are in contact with cattle and agricultural environments. While *Salmonella* is known to persist in the environment for long periods of time, this has primarily been documented in areas used for farming or areas where another source of contamination is present (Winfield and Groisman 2003, Toth et al. 2011). Another potential factor is the time of year during which samples were collected. Although we sampled birds from February through July, the two great-tailed grackles found to be positive for *Salmonella* Typhimurium were collected during the month of June. In cattle and horses, *Salmonella* shedding has been shown to follow distinct seasonal trends with summer and fall yielding the highest prevalence (Carter et al. 1986, Fossler et al. 2005, Cummings et al. 2009). It may be that *Salmonella* prevalence would increase if samples were collected during late summer and fall, as seen in livestock. As *Salmonella* thrives in warm, moist environments, this trend in seasonality is most likely related to temperature and moisture conditions that predominate in the summer and fall months (Strawn et al. 2013, Marine et al. 2015).

Resistance to antimicrobial agents that are included on the NARMS panel was not evident in the two study isolates. Presumably there is a lack of selection pressure for antimicrobial resistance among wild birds and other wildlife in most environments, relative to animals in agricultural settings. However, further research to generate a larger number of isolates would be necessary to evaluate the prevalence of antimicrobial resistance among *Salmonella* isolated from great-tailed grackles.

Both isolates were serotyped as *Salmonella* Typhimurium. According to the most recent CDC data, Typhimurium is the second most commonly reported serotype in human patients with laboratory-confirmed salmonellosis in the United States (Crim et al. 2015). This highlights the potential threat to public health posed by great-tailed grackles. Typhimurium is also the major cause of salmonellosis in wild birds and has been implicated in widespread outbreaks among songbirds in

the United States (Hernandez et al. 2012) and other areas of the world (Alley et al. 2002, Fukui et al. 2014). Mortality among infected birds can be high (Hernandez et al. 2012), and evidence suggests that the proportion of avian mortality attributed to salmonellosis may be increasing (Hall and Saito 2008). On a broader scale, Typhimurium is reported to be one of the most common serotypes among clinical and nonclinical laboratory-confirmed nonhuman sources (CDC 2014), underscoring the wide host range and complex ecology of this serotype.

### Acknowledgments

The authors would like to thank the students enrolled in Field and Lab Techniques in Vector-borne Disease Ecology (ENTO/VIBS 489-689) at Texas A&M University for their assistance with capturing and processing birds. The authors appreciate permission to capture birds from the parking lots of several businesses in College Station, Texas. This study was funded, in part, by teaching enhancement funds in the Department of Veterinary Integrative Biosciences, the Department of Entomology, and the Montague-Center for Teaching Excellence award to S.A.H.

### Author Disclosure Statement

No competing financial interests exist.

### References

- Alley MR, Connolly JH, Fenwick SG, Mackereth GF, et al. An epidemic of salmonellosis caused by *Salmonella* Typhimurium DT160 in wild birds and humans in New Zealand. *N Z Vet J* 2002; 50:170–176.
- Andrews W, Hammack T. *Bacteriological Analytical Manual: Salmonella*. Silver Spring, MD: U.S. Department of Health and Human Services, Food and Drug Administration (FDA), 2007.
- Callaway TR, Edrington TS, Nisbet DJ. Isolation of *Escherichia coli* O157: H7 and *Salmonella* from migratory brown-headed cowbirds (*Molothrus ater*), common grackles (*Quiscalus quiscula*), and cattle egrets (*Bubulcus ibis*). *Foodborne Pathog Dis* 2014; 11:791–794.
- Carter JD, Hird DW, Farver TB. Salmonellosis in hospitalized horses: Seasonality and case fatality rates. *J Am Vet Med Assoc* 1986; 188:163–167.
- Centers for Disease Control and Prevention (CDC). *National Salmonella Surveillance Annual Report, 2012*. Atlanta, GA: U.S. Department of Health and Human Services, CDC, 2014.
- Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals; Approved Standard—Third Edition*, Wayne, PA, 2008.
- Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement*, Wayne, PA, 2012.
- Crim SM, Griffin PM, Tauxe R, Marder EP, et al. Preliminary incidence and trends of infection with pathogens transmitted commonly through food—Foodborne Diseases Active Surveillance Network, 10 U.S. sites, 2006–2014. *MMWR Morb Mortal Wkly Rep* 2015; 64:495–499.
- Cummings KJ, Divers TJ, McDonough PL, Warnick LD. Fecal shedding of *Salmonella* spp among cattle admitted to a veterinary medical teaching hospital. *J Am Vet Med Assoc* 2009; 234:1578–1585.
- Fitzwater WD, Crabb AC, Marsh RE. Proceedings of the Vertebrate Pest Conference: Solutions to Urban Bird Problems. Davis, CA, 1988; 254–259.
- Food and Drug Administration. National Antimicrobial Resistance Monitoring System—Enteric Bacteria (NARMS): 2011 Executive Report. FDA, Rockville, MD, 2013.
- Fossler CP, Wells SJ, Kaneene JB, Ruegg PL, et al. Cattle and environmental sample-level factors associated with the presence of *Salmonella* in a multi-state study of conventional and organic dairy farms. *Prev Vet Med* 2005; 67:39–53.
- Fukui D, Takahashi K, Kubo M, Une Y, et al. Mass mortality of Eurasian Tree Sparrows (*Passer montanus*) from *Salmonella* Typhimurium DT40 in Japan, winter 2008–09. *J Wildl Dis* 2014; 50:484–495.
- Hall AJ, Saito EK. Avian wildlife mortality events due to salmonellosis in the United States, 1985–2004. *J Wildl Dis* 2008; 44:585–593.
- Hall DW, Harvey TM. Mortality at a night roost of Great-tailed Grackles and European Starlings during a spring hail storm. *Wilson J Ornithol* 2007; 119:309–312.
- Hamer SA, Lehrer E, Magle SB. Wild birds as sentinels for multiple zoonotic pathogens along an urban to rural gradient in greater Chicago, Illinois. *Zoonoses Public Health* 2012; 59: 355–364.
- Hernandez SM, Keel K, Sanchez S, Trees E, et al. Epidemiology of a *Salmonella enterica* subsp. *enterica* serovar Typhimurium strain associated with a songbird outbreak. *Appl Environ Microbiol* 2012; 78:7290–7298.
- Heymann DL. *Control of Communicable Diseases Manual: An Official Report of the American Public Health Association (APHA)*. Washington, DC: APHA Press, 2015.
- Janecko N, Čížek A, Halová D, Karpíšková R, et al. Prevalence, characterization and antibiotic resistance of *Salmonella* isolates in large corvid species of Europe and North America between 2010 and 2013. *Zoonoses Public Health* 2015; 62: 292–300.
- Kim JS, Lee GG, Park JS, Jung YH, et al. A novel multiplex PCR assay for rapid and simultaneous detection of five pathogenic bacteria: *Escherichia coli* O157: H7, *Salmonella*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Vibrio parahaemolyticus*. *J Food Prot* 2007; 70: 1656–1662.
- Kirk JH, Holmberg CA, Jeffrey JS. Prevalence of *Salmonella* spp. in selected birds captured on California dairies. *J Am Vet Med Assoc* 2002; 220:359–362.
- Marine SC, Pagadala S, Wang F, Pahl DM, et al. The growing season, but not the farming system, is a food safety risk determinant for leafy greens in the mid-Atlantic region of the United States. *Appl Environ Microbiol* 2015; 81: 2395–2407.
- Phalen DN, Drew ML, Simpson B, Roset K, et al. *Salmonella enterica* subsp. *enterica* in Cattle Egret (*Bubulcus ibis*) chicks from central Texas: Prevalence, serotypes, pathogenicity, and epizootic potential. *J Wildl Dis* 2010; 46:379–389.
- Rumbold DG, Morrison M, Bruner MC. Assessing the ecological risk of a municipal solid waste landfill to surrounding wildlife: A case study in Florida. *Environ Bioindicators* 2009; 4:246–279.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 2011; 17:7–15.
- Scharff RL. Economic burden from health losses due to foodborne illness in the United States. *J Food Prot* 2012; 75:123–131.

- Skov MN, Madsen JJ, Rahbek C, Lodal J, et al. Transmission of *Salmonella* between wildlife and meat-production animals in Denmark. *J Appl Microbiol* 2008; 105:1558–1568.
- Strawn LK, Fortes ED, Bihn EA, Nightingale KK, et al. Landscape and meteorological factors affecting prevalence of three food-borne pathogens in fruit and vegetable farms. *Appl Environ Microbiol* 2013; 79:588–600.
- Toth JD, Aceto HW, Rankin SC, Dou Z. Survival characteristics of *Salmonella enterica* serovar Newport in the dairy farm environment. *J Dairy Sci* 2011; 94:5238–5246.
- Wehtje W. The range expansion of the Great-tailed Grackle (*Quiscalus mexicanus* Gmelin) in North America since 1880. *J Biogeogr* 2003; 30:1593–1607.
- Winfield MD, Groisman EA. Role of nonhost environments in the lifestyles of *Salmonella* and *Escherichia coli*. *Appl Environ Microbiol* 2003; 69:3687–3694.

Address correspondence to:

Kevin J. Cummings

Department of Veterinary Integrative Biosciences

College of Veterinary Medicine & Biomedical Sciences

Texas A&M University

College Station, TX 77843-4458

E-mail: kcummings@cvm.tamu.edu