

About the Author

Dr. de Sousa is a public health microbiologist responsible for the Rickettsial and the Gastrointestinal Viral Infections Units at the National Institute of Health Dr. Ricardo Jorge. She works on research and diagnosis of rickettsial diseases.

References

- Jado I, Oteo JA, Aldámiz M, Gil H, Escudero R, Ibarra V, et al. *Rickettsia monacensis* and human disease, Spain. *Emerg Infect Dis*. 2007;13:1405–7. <https://doi.org/10.3201/eid1309.060186>
- Madeddu G, Mancini F, Caddeo A, Ciervo A, Babudieri S, Maida I, et al. *Rickettsia monacensis* as cause of Mediterranean spotted fever-like illness, Italy. *Emerg Infect Dis*. 2012;18:702–4. <https://doi.org/10.3201/eid1804.111583>
- Kim YS, Choi YJ, Lee KM, Ahn KJ, Kim HC, Klein T, et al. First isolation of *Rickettsia monacensis* from a patient in South Korea. *Microbiol Immunol*. 2017;61:258–63. <https://doi.org/10.1111/1348-0421.12496>
- Kim SW, Kim CM, Kim DM, Yun NR. Case report: coinfection with *Rickettsia monacensis* and *Oraientia tsutsugamushi*. *Am J Trop Med Hyg*. 2019;101:332–5. <https://doi.org/10.4269/ajtmh.18-0631>
- Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, et al. Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev*. 2013;26:657–702. <https://doi.org/10.1128/CMR.00032-13>
- de Sousa R, Pereira BI, Nazareth C, Cabral S, Ventura C, Crespo P, et al. *Rickettsia slovaca* infection in humans, Portugal. *Emerg Infect Dis*. 2013;19:1627–9. <https://doi.org/10.3201/eid1910.130376>
- de Sousa R, Barata C, Vitorino L, Santos-Silva M, Carrapato C, Torgal J, et al. *Rickettsia sibirica* isolation from a patient and detection in ticks, Portugal. *Emerg Infect Dis*. 2006;12:1103–8. <https://doi.org/10.3201/eid1207.051494>
- Sousa R, França A, Dória Nóbrega S, Belo A, Amaro M, Abreu T, et al. Host- and microbe-related risk factors for and pathophysiology of fatal *Rickettsia conorii* infection in Portuguese patients. *J Infect Dis*. 2008;198:576–85. <https://doi.org/10.1086/590211>
- Portillo A, de Sousa R, Santibáñez S, Duarte A, Edouard S, Fonseca IP, et al. Guidelines for the detection of *Rickettsia* spp. *Vector Borne Zoonotic Dis*. 2017;17:23–32. <https://doi.org/10.1089/vbz.2016.1966>
- Tijssse-Klasen E, Sprong H, Pandak N. Co-infection of *Borrelia burgdorferi* sensu lato and *Rickettsia* species in ticks and in an erythema migrans patient. *Parasit Vectors*. 2013;6:347.

Address for correspondence: Rita de Sousa, Av. Da Liberdade n°5, 2965, Águas de Moura, Portugal; email: rsr.desousa@gmail.com

Domestic Dogs as Sentinels for West Nile Virus but not *Aedes*-borne Flaviviruses, Mexico

Edward Davila,¹ Nadia A. Fernández-Santos,¹ José Guillermo Estrada-Franco, Lihua Wei, Jesús A. Aguilar-Durán, María de J. López-López, Roberto Solís-Hernández, Rosario García-Miranda, Doireyner Daniel Velázquez-Ramírez, Jasiel Torres-Romero, Susana Arellano Chávez, Raúl Cruz-Cadena, Roberto Navarro-López, Adalberto A. Pérez de León,² Carlos Guichard-Romero, Estelle Martin,³ Wendy Tang, Matthias Frank, Monica Borucki, Michael J. Turell, Alex Pauvolid-Corrêa, Mario A. Rodríguez-Pérez, Héctor Ochoa-Díaz-López, Sarah A. Hamer, Gabriel L. Hamer

Author affiliations: Texas A&M University, College Station, Texas, USA (E. Davila, E. Martin, W. Tang, A. Pauvolid-Corrêa, S.A. Hamer, G.L. Hamer); Instituto Politécnico Nacional, Reynosa, México (N.A. Fernández-Santos, J.G. Estrada-Franco, L. Wei, J.A. Aguilar-Durán, M. de J. López-López, M.A. Rodríguez-Pérez); El Colegio de la Frontera Sur, San Cristóbal de Las Casas, México (R. Solís-Hernández, R. García-Miranda, D.D. Velázquez-Ramírez, J. Torres-Romero, H. Ochoa-Díaz-López); Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, México (S. Arellano Chávez); Universidad Autónoma de Chiapas, Ocozocoautla de Espinosa, México (R. Cruz-Cadena); Comisión México-Estados Unidos para la Prevención de la Fiebre Aftosa y Otras Enfermedades Exóticas de los Animales, México City (R. Navarro-López); US Department of Agriculture Agricultural Research Service Knippling-Bushland Livestock Insects Research Laboratory, Kerrville, Texas, USA (A.A. Pérez de León); Zoológico Miguel Álvarez del Toro, Tuxtla Gutiérrez (C. Guichard-Romero); Lawrence Livermore National Laboratory, Livermore, California, USA (M. Frank, M. Borucki); VectorID LLC, Frederick, Maryland, USA (M.J. Turell)

DOI: <https://doi.org/10.3201/eid2805.211879>

We tested 294 domestic pet dogs in Mexico for neutralizing antibodies for mosquito-borne flaviviruses. We found high (42.6%) exposure to West Nile virus in Reynosa (northern Mexico) and low (1.2%) exposure in Tuxtla Gutiérrez (southern Mexico) but very limited exposure to *Aedes*-borne flaviviruses. Domestic dogs may be useful sentinels for West Nile virus.

¹These authors contributed equally to this article.

²Current affiliation: US Department of Agriculture Agricultural Research Service San Joaquin Valley Agricultural Sciences Center, Parlier, California, USA.

³Current affiliation: University of Florida, Gainesville, Florida, USA.

Mosquito-transmitted viruses represent substantial health burdens across the Americas. Despite the broad geographic ranges of *Aedes* spp. and *Culex* spp. mosquitoes, the endemicity of human arboviral diseases is incongruent with these vector distributions (1,2). Animal sentinels may therefore be useful for signaling areas of virus transmission and human risk, especially in resource-poor settings where human diseases may be underreported. Although *Ae. aegypti* mosquitoes have been considered to feed predominantly on humans and *Cx. quinquefasciatus* mosquitoes on birds, our recent work studying host feeding patterns in southern Texas, USA (3), and northern Mexico (4) has documented substantial feeding on dogs for both species, presenting a novel opportunity to evaluate dogs for possible sentinel surveillance. Because dogs are ubiquitous and share the domestic environment with humans, tracking their exposures might provide evidence for understanding human risk and a sensitive indicator of geographic variation for mosquito-borne disease risk. We aimed to estimate domestic dog exposure to Zika virus (ZIKV), dengue virus 1 (DENV-1) and DENV-2, and West Nile virus (WNV) in northern and southern Mexico based on the presence and quantity of specific neutralizing antibodies as a proxy for human risk.

During 2018–2019, we sampled pet dogs from 3 residential areas in the city of Tuxtla Gutierrez, Chiapas, in southern Mexico and 8 neighborhoods in

the city of Reynosa, Tamaulipas, in northern Mexico (Figure). We initially screened serum or plasma samples at a 1:10 dilution, then further tested those that neutralized PFUs by $\geq 90\%$ in duplicates at serial 2-fold dilutions that ranged from 1:10 to 1:320 to determine 90% endpoint titers (Appendix, <https://wwwnc.cdc.gov/EID/article/28/5/21-1879-App1.pdf>).

We tested blood samples from 294 pet dogs (predominantly mixed breeds, chihuahuas, and pit bulls). Canine exposure to WNV was widespread, and we found a higher prevalence of neutralizing antibodies to WNV in dogs from Reynosa (72/169, 42.6%) than in those from Tuxtla Gutierrez (1/87, 1.2%; Appendix). In contrast, only 2 (0.7%) dogs from Tuxtla Gutierrez had neutralizing antibodies for ZIKV exposure, showing endpoint titers of 40 and 10. However, the dog with a ZIKV titer of 40 also had a 90% plaque-reduction neutralization test titer of 20 for WNV; we could not screen the dog with a ZIKV titer of 10 for other viruses because of low sample volume. A single dog from Tuxtla Gutierrez had a low titer monotypic reaction for DENV-2, the only evidence of exposure to an *Aedes*-borne flavivirus (Appendix). A sample size analysis indicated that the level of sampling we conducted supports 95% confidence that true prevalence of neutralizing antibodies in these canine populations did not exceed 1% for each of these *Aedes*-borne flaviviruses.




Figure. Sampling locations in Tuxtla Gutierrez, Chiapas, and Reynosa, Tamaulipas, Mexico, for study of neutralizing antibodies for mosquito-borne flaviviruses in domestic dogs. Map was created using QGIS 3.18.2 (<https://qgis.org/en/site>) with public domain map data from Instituto Nacional de Estadística, Geografía e Informática (National Institute of Statistics, Geography, and Computer Science [INEGI]; <https://www.inegi.org.mx/app/mapas>) and satellite images from Google Maps (<https://www.google.com.mx/maps>).

Our data suggested substantial WNV enzootic activity in Reynosa and corroborated prior observations of high use of dogs as blood meal hosts by *Cx. quinquefasciatus* mosquitoes. Despite detecting neutralizing antibodies for WNV in 42.6% of dogs from Reynosa, the number of reported human WNV cases in Mexico has remained low (5), suggesting that transmission occurs among domestic animals but either humans have not been infected or cases have not been reported. Texas has a high number of reported human WNV cases (Texas Department of State Health Services, <https://dshs.texas.gov/idcu/disease/arboviral/westNile/#stats>). The lower reported numbers of WNV cases in Mexico might be in part because of the high seroprevalence of antibodies for other flaviviruses, which have been shown to protect against severe clinical infection from WNV, thus leading to reduced testing (6). Low WNV seroprevalence among dogs in Tuxtla Gutierrez might reflect a larger diversity of vertebrates with lower WNV competence, fed upon by *Culex* mosquitoes in the study area.

The relative lack of canine exposure to *Aedes*-borne flaviviruses suggests not an absence of these viruses circulating in these communities but that dogs are likely insensitive sentinels of the viruses' transmission in Mexico. In Chiapas, 7,972 human cases of dengue and 763 cases of Zika had been reported during 2016–2020 (7,8). Considering the timing of our sampling and the ages of the dogs, we expect that ≈75% of sampled dogs were living in these communities during DENV and ZIKV transmission activity. In the state of Tamaulipas, there were 3,988 human cases of dengue (7) and 733 cases of Zika during 2016–2020 (8). Given recent quantification that >50% of *Ae. aegypti* in southern Texas and northern Mexico feed on dogs (3,4), our serologic data suggest that either the probability of virus spillover into dogs is low or that, although dogs are susceptible to infection, neutralizing antibodies developed weakly or waned rapidly (9).

Our study suggests substantial WNV enzootic activity in Reynosa, Mexico and corroborates observations that *Cx. quinquefasciatus* mosquitoes, a primary vector of WNV, use high numbers of dogs for blood meals. Therefore, domestic pet dogs may be useful sentinels of WNV transmission, as previously suggested in other regions (10).

Acknowledgments

We thank the World Reference Center for Emerging Viruses and Arboviruses at the University of Texas Medical Branch and the Centers for Disease Control and Prevention for providing the viruses used in this study.

We appreciate field sampling assistance in Tamaulipas from Sofia Rodríguez, Irma Cobos, Cristian Delgado, Mónica Duarte, Diana Navarrate, Elisa Rodarte, Luis Sánchez, Ricardo Palacios, Adebisi Adeniran, and Ester Carbajal. We appreciate field sampling assistance in Chiapas from Paola Ruiz, Daniela Mendoza, Ali Fajardo, Azucena, Katia Hernandez, Ma. Fernanda Escobar, Emiliano Escobar, Nathan Penagos, and Cristel Nandayapa.

Our work was performed, in part, under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344 to G.L.H., M.F., M.K.B. Additional support came from the Texas A&M University-Consejo Nacional de Ciencia y Tecnología Collaborative Research Program grant (no. 2018-041-1) and a Texas A&M AgriLife Insect Vector seed grant. J.G.E.F. was supported by grants from Secretaría de Investigación y Posgrado of Instituto Politécnico Nacional (Nos. 20196759, 20200843, and 20202442).

About the Author

Mr. Davila is a PhD student at the Texas A&M University College of Veterinary Medicine and Biomedical Sciences. His primary research interests include animals as sentinels for human disease and the epidemiology of emerging infectious diseases.

References

1. Kraemer MU, Sinka ME, Duda KA, Mylne AQ, Shearer FM, Barker CM, et al. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *eLife*. 2015;4:e08347. <https://doi.org/10.7554/eLife.08347>
2. Rochlin I, Faraji A, Healy K, Andreadis TG. West Nile virus mosquito vectors in North America. *J Med Entomol*. 2019;56:1475–90. <https://doi.org/10.1093/jme/tjz146>
3. Olson MF, Ndeffo-Mbah ML, Juarez JG, Garcia-Luna S, Martin E, Borucki MK, et al. High rate of non-human feeding by *Aedes aegypti* reduces Zika virus transmission in South Texas. *Viruses*. 2020;12:E453. <https://doi.org/10.3390/v12040453>
4. Estrada-Franco JG, Fernández-Santos NA, Adebisi AA, López-López MJ, Aguilar-Durán JA, Hernández-Triana LM, et al. Vertebrate-*Aedes aegypti* and *Culex quinquefasciatus* (Diptera) arbovirus transmission networks: non-human feeding revealed by meta-barcoding and next-generation sequencing. *PLoS Negl Trop Dis*. 2020;14:e0008867. <https://doi.org/10.1371/journal.pntd.0008867>
5. Elizondo-Quiroga D, Elizondo-Quiroga A. West Nile virus and its theories, a big puzzle in Mexico and Latin America. *J Glob Infect Dis*. 2013;5:168–75. <https://doi.org/10.4103/0974-777X.122014>
6. Tesh RB, Travassos da Rosa AP, Guzman H, Araujo TP, Xiao SY. Immunization with heterologous flaviviruses protective against fatal West Nile encephalitis. *Emerg Infect Dis*. 2002;8:245–51. <https://doi.org/10.3201/eid0803.010238>
7. Dirección General de Epidemiología, Secretaría de Salud México. Historical epidemiological bulletin [in Spanish]

- [cited 2022 Jan 26]. <https://www.gob.mx/salud/acciones-y-programas/historico-boletin-epidemiologico>
8. Pan American Health Organization. Zika: confirmed cases in Mexico, 2017 [cited 2020 Jun 19]. https://www3.paho.org/data/index.php/en/?option=com_content&view=article&id=526:zika-mex-en&Itemid=352
 9. Thongyuan S, Kittayapong P. First evidence of dengue infection in domestic dogs living in different ecological settings in Thailand. *PLoS One*. 2017;12:e0180013. <https://doi.org/10.1371/journal.pone.0180013>
 10. Resnick MP, Grunenwald P, Blackmar D, Hailey C, Bueno R, Murray KO. Juvenile dogs as potential sentinels for West Nile virus surveillance. *Zoonoses Public Health*. 2008;55:443–7. <https://doi.org/10.1111/j.1863-2378.2008.01116.x>

Address for correspondence: Gabriel Hamer, Texas A&M University, TAMU 2475, College Station, TX, 77843, USA; email: gghamer@tamu.edu; Héctor Ochoa-Díaz-López, El Colegio de la Frontera Sur, Departamento de Salud, San Cristóbal de Las Casas, Chiapas, México, CP. 29290; email: hochoa@ecosur.mx; Mario A. Rodríguez-Pérez, Instituto Politécnico Nacional, Centro de Biotecnología Genómica, Blvd. del Maestro esquina Elías Piña s/n, Colonia Narciso Mendoza, 88170, Cd. Reynosa, Tamaulipas, México; email: mrodriguez@ipn.mx

Viral Hepatitis E Outbreaks in Refugees and Internally Displaced Populations, sub-Saharan Africa, 2010–2020

Angel N. Desai, Amir M. Mohareb, Mubarak Mustafa Elkarsany, Hailemichael Desalegn, Lawrence C. Madoff, Britta Lassmann

Author affiliations: University of California–Davis, Sacramento, California, USA (A.N. Desai); International Society for Infectious Diseases, Brookline, Massachusetts, USA (A.N. Desai, M.M. Elkarsany, L.C. Madoff, B. Lassmann); Massachusetts General Hospital, Boston, Massachusetts, USA (A.M. Mohareb); Karary University, Khartoum, Sudan (M.M. Elkarsany); St. Paul's Hospital MMC, Addis Ababa, Ethiopia (H.D. Desalegn); University of Massachusetts Medical School, Worcester, Massachusetts, USA (L.C. Madoff)

DOI: <https://doi.org/10.3201/eid2805.212546>

Hepatitis E virus is a common cause of acute viral hepatitis. We analyzed reports of hepatitis E outbreaks among forcibly displaced populations in sub-Saharan Africa during 2010–2020. Twelve independent outbreaks occurred, and >30,000 cases were reported. Transmission was attributed to poor sanitation and overcrowding.

Hepatitis E virus (HEV) is a common etiology of acute viral hepatitis worldwide (1). Large-scale, often protracted outbreaks caused by HEV infection in refugee and internally displaced person (IDP) settlements and camps have occurred (1), particularly in sub-Saharan Africa, a region with nearly one third of the global forcibly displaced population (2). Previous epidemiologic studies of HEV infections in forcibly displaced persons have focused on singular events (3,4). The objective of this study was to identify trends in HEV outbreaks among forcibly displaced populations in sub-Saharan Africa.

We conducted a focused review of all English-language curated reports posted on ProMED-mail (ProMED) during 2010–2020 concerning HEV in forcibly displaced populations in sub-Saharan Africa. ProMED uses formal and informal disease surveillance mechanisms to rapidly report emerging disease events in animals, humans, and plants globally (5). It has been validated as a rapid and accurate tool for determining and describing global outbreaks. We verified all reports via PubMed, ReliefWeb, the UN High Commission for Refugees, World Health Organization (WHO), and references secondarily collected from ProMED. We used the keyword “hepatitis E” in applicable search engines for reports published during 2010–2020. We included records documenting “refugee(s) and/or asylum seeker(s) and/or internally displaced person(s)” in sub-Saharan Africa as defined by the World Bank (6). We considered outbreaks unique on the basis of date and location of cases. When screening ProMED reports, we used the most recent report pertaining to an outbreak. In cases where discrepancies existed between data sources reporting on the same outbreak, we retained the higher number of case counts. Three independent investigators (A.D., B.L., and A.M.) manually reviewed the databases.

Twelve hepatitis E outbreaks among forcibly displaced persons resulting in a total of >30,000 suspected or confirmed cases of acute HEV and ≥610 deaths were reported during 2010–2020 (Appendix Table, <https://wwwnc.cdc.gov/EID/article/28/5/21-2546-App1.pdf>). Outbreaks occurred in Sudan, South Sudan, Ethiopia, Chad, Niger, Namibia, Burkina Faso, Kenya, and Nigeria (Figure). One outbreak in displaced persons in South Sudan’s Bentiu camp for